

## Supplementary file 1

**Table S1.** Co-treatment of PTX and SNPs against MDA-MB 231: A) Experimental range and level of SNPs and PTX concentrations. B) RSM to study the synergistic effect of SNPs and PTX on MDA-MB 231 cell line viability. C) Analysis of variance for response surface 2FI Model. D) Optimum values of SNPs and PTX in coded and none-coded forms.

Variables	Symbol coded	Range and level		
		-1	0	+1
paclitaxel (μM)	X <sub>1</sub>	0.01	1.505	3
SNP (μM)	X <sub>2</sub>	1	4	7

Comment	X <sub>1</sub>	X <sub>2</sub>	Cell viability (%)
Center	0	0	52
Center	0	0	51
Center	0	0	53
Center	0	0	52
Center	0	0	52
Axial	0	-1.02	58
Axial	-1.02	0	66
Axial	1.02	0	47
Axial	0	1.02	46
Fact	1	-1	50
Fact	1	1	33
Fact	-1	-1	100
Fact	-1	1	50

Source	Sum of Squares	df	Mean Square	F value	p-value
X <sub>1</sub>	1230.998	1	1230.998	32.784	0.0003
X <sub>2</sub>	1037.899	1	1037.899	27.642	0.0005
X <sub>1</sub> X <sub>2</sub>	272.250	1	272.250	7.250	0.0247

Variable	X <sub>1</sub>	X <sub>2</sub>	Cell viability
Coded	-0.318	0.583	50
Non coded	1.03 (μM)	5.75 (μM)	50

The obtained data in B, last column, were analyzed by Design Expert Software. The coefficients of a regression equation were calculated with R<sup>2</sup> of 0.88:

$$Y = 97.58090 - 16.91972 X_1 - 7.14373 X_2 + 1.83946 X_1 X_2$$

Where, Y is the percent of MDA-MB 231 cells viability, X<sub>1</sub> and X<sub>2</sub> are the concentration (μM) of PTX and SNPs, respectively. According to this equation, in addition to the linear effects of X<sub>1</sub> and X<sub>2</sub>, the interaction effect of X<sub>1</sub> and X<sub>2</sub> is also effective on cell viability (P < 0.05).



**Table S2.** Optimization of SNPs-PTX@BSANPs preparation. A) Experimental range and level of BSA, BSA-Drug and pH, B) Experimental design through the RSM to study optimization of the variables based on NPs size , C) Analysis of variance for Response Surface-Reduced Quadratic Model. D) Optimum values of BSA, BSA-Drug and pH in coded and none coded forms.


Variables	Symbol coded	Range and level		
		-1	0	+1
BSA (%)	A	1	3	5
BSA:Drug ( $\mu\text{M}$ : $\mu\text{M}$ )	B	1.5	5.25	9
pH	C	5.40	6.40	7.40

Source	Sum of Squares	df	Mean Square	F value	p-value
A	7460.547	1	7460.547	603.3909	< 0.0001
B	1618.016	1	1618.016	130.8612	< 0.0001
A <sup>2</sup>	2977.024	1	2977.024	240.7745	< 0.0001
B <sup>2</sup>	1827.456	1	1827.456	147.8002	< 0.0001


Variable	A	B	C	Size
Coded	-0.79	+0.315	+0.88	50
Non coded	1.42	6.43	7.28	50

Comment	BSA% (A)	BSA: SNPs-PTX@BSANPs (B)	pH (C)	Size (nm)
Center	0	0	0	58
Center	0	0	0	58
Center	0	0	0	58.1
Center	0	0	0	58.1
Center	0	0	0	58
Center	0	0	0	58.6
Axial	0	-1.3	0	106.4
Axial	0	0	-1.3	58
Axial	0	1.3	0	70
Axial	0	0	1.3	60
Axial	-1.3	0	0	60
Axial	1.3	0	0	131.7
Fact	-1	1	1	50.9
Fact	-1	-1	1	80.1
Fact	1	-1	1	130
Fact	1	1	-1	105
Fact	1	-1	-1	130
Fact	-1	1	-1	65.91
Fact	1	1	1	110
Fact	-1	-1	-1	80


The obtained data in B-last column, were analyzed by design expert software. The coefficients of the regression equation were calculated according to the following equation with  $R^2$  of 0.98:  
 $Z = 114.07771 - 17.84403 A - 15.10180 B + 5.09986 A^2 + 1.13655B^2$   
 Where, Z is the size of NPs, A is the percent of BSA and B is the ratio of BSA to anticancer agents. In addition to A and B, the square of A and B also affect the size of NPs ( $P < 0.05$ ). However, the effect of pH was not significant..




**Figure S1.** Two-dimensional contour plots of cell viability of MDA-MB 231 ( $1 \times 10^4$  cell/ml) treated with different concentrations of PTX (0.01 to 3  $\mu$ M) and SNPs (1 to 7  $\mu$ M) simultaneously. The sign  $\bullet$  designate the designed points. The color change from red to blue indicates reducing the cell viability.



**Figure S2.** Correlation between the calculated and experimental values of cell viability in synergistic system of PTX and SNPs. The color change from red to blue indicates reducing the cell viability.



**Figure S3.** Two-dimensional contour plots of the effect of BSA percent (horizontal axis) and ratio of BSA: Drug (Vertical axis) on size of SNPs-PTX@BSANPs. The color change from red to blue indicates reducing the diameter from 131.7 to 50.9 nm.



**Figure S4.** Correlation between calculated and experimental values of size of NPs in optimization of SNPs-PTX@BSANPs preparation. The color change from red to blue indicates reducing the size of NPs.